



**MAKHTESHIM**  
**A G A N**  
NORTH AMERICA



# Arrow™ 2EC HERBICIDE

## ACTIVE INGREDIENT

% BY WT.

Clethodim: (E)-2-[1-[(3-chloro-2-propenyl)oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one	26.4%
<b>OTHER INGREDIENTS:</b>	73.6%
<b>TOTAL</b>	100.0%

Contains petroleum distillate.

Contains 2.0 lbs. clethodim per gallon.

EPA Reg. No. 66222-60 EPA Est. No. 5905-GA-001<sup>HC</sup>  
5905-IA-001<sup>HD</sup>

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

**FOR EMERGENCY MEDICAL HELP, CALL PROSAR AT 1-877-250-9291, 24 HOURS.**

For additional precautionary, handling and use statements, and First Aid instructions see inside of this booklet.

Letter in lot number corresponds to superscript in EPA Est. No.

This product is not yet registered in California.



**Makhteshim Agan of North America, Inc.**

**MANA**

551 Fifth Avenue / Suite 1100  
New York, NY 10176

**MAKHTESHIM**  
**A G A N**  
NORTH AMERICA

**Net Contents: 1 gallon**

# KEEP OUT OF REACH OF CHILDREN CAUTION

## FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.
- IF ON SKIN OR CLOTHING:**
- Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15-20 minutes.
  - Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:**
- Immediately call a poison control center or doctor.
  - Do not induce vomiting unless told to do so by a poison control center or doctor.
  - Do not give any liquid to the person.
  - Do not give anything by mouth to an unconscious person.
- IF INHALED:**
- Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
  - Call a poison control center or doctor for further treatment advice.

**Have the product container or label with you when calling a poison control center or doctor or going for treatment.** You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. Contains petroleum distillate.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category **G** on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or Viton  $\geq 14$  mils
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwater or rinsate.

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist:

Solano Grass:	Solano County, California: the vernal lakes area bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south, and Travis Air Force Base to the west.
Wild Rice:	Hays County, Texas

## PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for

*(continued)*

### **AGRICULTURAL USE REQUIREMENTS *(continued)***

training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or Viton  $\geq$  14 mils
- Shoes plus socks

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of operating areas, or vicinity where there may be drift. Do not enter treated area without protective clothing until sprays have dried.

### **TANK MIXES**

**NOTICE:** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator, and/or application advisor. Read and follow the entire label of each product to be used in the tank mix with this product.

### **GENERAL INFORMATION**

**FOR USE ON:** Soybeans, Cotton, Ornamentals, Sugarbeets, Onions (dry bulbs and green), Garlic, Shallots (dry bulbs and green), Alfalfa, Peanuts, Dry Beans,

Sunflower, Canola, Flax, Mustard Seed, Potato, Sweet Potato, Yam (and other Tuberous<sup>1</sup> and Corm<sup>1</sup> Vegetables), Tomatoes, Peppers (bell and non-bell), Eggplants (and other Fruiting Vegetables), Carrot, Radish, Garden Beet, Horseradish (and other Root Vegetables<sup>2</sup>), Leaf Lettuce, Broccoli, Cabbage, Cauliflower (and other Head and Stem Brassica Vegetables<sup>3</sup>), Mustard Greens (and other Leafy Brassica Greens<sup>4</sup>), Spinach, Celery, Rhubarb (and other Leaf Petioles<sup>5</sup>), Cranberry, Strawberry, Squash (including Pumpkins), Cucumber, Melons (including Cantaloupes and Watermelons), Mint, Clover (grown in Idaho, Oregon, and Washington only), Conifer Trees, Non-Bearing Food Crops, Fallow Land (and other non-producing agricultural areas), and Non-crop or Non-planted Areas.

<sup>1</sup> Other tuber and corm vegetables approved for use with Arrow 2EC Herbicide include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible burdock, edible canna, bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, tanier, turmeric, and bean yam.

<sup>2</sup> Other root vegetables approved for use with Arrow 2EC Herbicide include: burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret, and turnip.

<sup>3</sup> Other head and stem brassica vegetables approved include: Chinese broccoli; Brussels sprouts; Chinese (Napa) cabbage; Chinese mustard; cavalo broccolo; and kohlrabi.

<sup>4</sup> Other leafy brassica greens approved for use with Arrow 2EC Herbicide include: broccoli raab, cabbage, Chinese (Bok Choy), collards, kale, mizuna, mustard greens, mustard spinach, rape greens, and turnip greens.

<sup>5</sup> Other leaf petiole crops include: cardoon, Chinese celery, celtuce, Florence fennel, and swiss chard.

Arrow 2EC Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

Arrow 2EC Herbicide is a selective postemergence herbicide for control of annual and perennial grasses. Arrow 2EC Herbicide does not control sedges or broadleaf weeds.

Repeated use of Arrow 2EC Herbicide (or similar postemergence grass herbicide with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species.

If poor performance occurs and cannot be attributed to adverse weather or application conditions, a resistant biotype may be present. This is most likely to occur in fields where other control strategies such as crop rotation, mechanical removal, and other classes of herbicides are not used from year to year.

Do not allow Arrow 2EC Herbicide to come in contact with desirable grass crops such as corn, rice, sorghum, small grains, or turf, as these and other grass crops will be injured or killed. Minor leaf spotting may occur on treated plants under certain environmental conditions. New foliage is not affected.

#### **Control symptoms**

Treated grass weeds show a reduction in vigor and growth. Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days depending on grass species treated and environmental conditions.

## **APPLICATION INFORMATION**

#### **Timing of Applications**

Apply Arrow 2EC Herbicide postemergence to actively growing grasses according to rate table recommendations. Applications made to grass plants stressed by insufficient moisture or cold temperatures or to grass plants exceeding recommended growth stages may result in unsatisfactory control. Do not apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, Arrow 2EC Herbicide should be applied as soon as possible after an irrigation (within 7 days). In arid regions, a second application of Arrow 2EC Herbicide will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of Arrow 2EC Herbicide may reduce weed control. DO NOT APPLY Arrow 2EC Herbicide if rainfall is expected within one hour since control may be reduced.

## ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE

<b>CROP</b>	<b>ADJUVANT RECOMMENDATIONS</b>
Soybeans, Alfalfa, Dry Beans, Cotton, Peanuts, Sugar Beets, Sunflowers, Potatoes	Always use a crop oil concentrate* at 1.0 qt./A by ground or 1% v/v (but not less than 1 pt./A) in the finished spray volume by air.  1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to Arrow 2EC Herbicide applications in addition to the recommended rate of crop oil concentrate. The addition of AMS has shown improved grass control for difficult to control species including, quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
Onions (dry bulbs and green), Garlic, Shallots (dry bulbs and green), Carrots, Radishes, Garden Beets, Horseradish (and other Root Vegetables), Leaf Lettuce, Broccoli, Cabbage, Cauliflower (and other Head and Stem Brassica Vegetables), Mustard Greens (and other Leafy Brassica Greens), Spinach, Celery, Rhubarb (and other Leaf Petioles), Cranberries, Sweet Potatoes, Yams (and other tuberous and corm vegetables),	Always use a crop oil concentrate at 1% v/v in the finished spray volume unless tank mix instructions indicate otherwise.  Addition of liquid fertilizer is not recommended for these crops.



CROP	ADJUVANT RECOMMENDATIONS
Canola, Flax, Mustard Seed, Tomatoes, Peppers (bell and non-bell), Eggplants (and other fruiting vegetables), Strawberries, Squash (including Pumpkins), Cucumbers, Melons (including Cantaloupes and Watermelons), Mint, Clover	Always use a crop oil concentrate at 1% v/v in the finished spray volume unless tank mix instructions indicate otherwise. Addition of liquid fertilizer is not recommended for these crops.
Ornamental Plants, Non-Bearing Food Crops	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v). Use of crop oil concentrate is not recommended since it may injure flowers and foliage.
Conifer Trees, Fallow Land (and other non-producing agricultural areas), and Non-Crop or Non-Planted Areas	Always use a crop oil concentrate containing at least 15% emulsifier at 1% v/v (but not less than 1 pt./A) in the finished spray volume.

\*Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality, and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

#### **Ground Application**

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 40 gals. of spray solution per acre. Under the following conditions a minimum of 10 gals. per acre is required: narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions, heavy grass pressure, or

when grasses are at or near maximum height. Failure to use a minimum of 10 gals. per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Applications to onions (dry bulbs and green), garlic, or shallots (dry bulbs and green) should be made in a minimum of 20 gals. of spray solution per acre.

#### **Air application**

Use a minimum of 3 gals. of spray solution per acre. Increase spray volumes up to 10 gals. as grass or crop foliage becomes dense.

For onions (dry bulbs and green), garlic, or shallots (dry bulbs and green): When applying by air do not exceed 8 fl. oz./A in a single application.

**Note: Crop injury may occur when Arrow 2EC Herbicide is applied to onions, garlic, or shallots with aerial equipment.**

#### **Spot Treatment**

When using hand sprayers or high volume sprayers utilizing hand guns, mix 1/4% to 1/2% (0.33 oz. to 0.65 oz. per gal.) Arrow 2EC Herbicide and treat to wet vegetation while not allowing runoff of spray solution. For uses requiring crop oil concentrate, include crop oil concentrate at 1% (1.3 oz. per gal.) by volume. For uses requiring non-ionic surfactant, include non-ionic surfactant at 1/4% (0.33 oz. per gal.) by volume.

**Note: If Arrow 2EC Herbicide is applied as a spot treatment, care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.**

## **CHEMIGATION ONIONS (Dry Bulbs and Green) AND GARLIC SPRINKLER IRRIGATION APPLICATION**

This product may be applied to onions and garlic by sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set, or hand move. Do not apply by chemigation to any other crop or to this crop using any other type of irrigation system.

Apply Arrow 2EC Herbicide at the high rate recommended for annual grasses (16 fl. oz./A) when the grass height is at the low end of the range (application to larger grasses may not provide adequate control). Add a crop oil concentrate containing at least 15% emulsifier at 1 quart per acre.

Apply Arrow 2EC Herbicide in 0.1 to 0.2 acre-inch of water either at the end of a regular irrigation set or as a separate application not associated with a regular irrigation using the least amount of water that provides proper distribution and coverage. Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness. Use a metering device to inject the Arrow 2EC Herbicide into the irrigation water at a constant flow. Constant agitation must be maintained in the chemical supply tank during the entire period of herbicide application. Inject the product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period.

It is not recommended that Arrow 2EC Herbicide be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

### **Use Precautions**

1. Apply this product only through irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
3. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

5. A person knowledgeable of the chemigation system and responsible for its operation or under supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
12. Do not apply when wind speed favors drift beyond the area intended for treatment.

## **RESTRICTIONS AND LIMITATIONS**

### **GENERAL**

Do not apply if rain is expected within 1 hour of application as control may be unsatisfactory.

Do not apply a postemergence broadleaf herbicide within one day following application of Arrow 2EC Herbicide or reduced grass control may result.

Arrow 2EC Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

For canola, flax, and mustard seed crops, do not apply more than 5 fl. oz. of Arrow 2EC Herbicide (0.08 lb.ai) per acre per season. For clover and radish crops, do not apply more than 16 fl. oz. of Arrow 2EC Herbicide (0.25 lb. ai) per acre per season. For all other crops, do not apply more than 32 fl. oz. of Arrow 2EC Herbicide (0.50 lb. ai) per acre per season. Application on Long Island, New York is restricted to no more than 16 fl. oz. of Arrow 2EC Herbicide (0.25 lb.ai) per acre per season.

Do not apply more than 8 fl. oz./A of Arrow 2EC Herbicide **per application** to the following crops: garden beets, carrots, radish (and other root vegetables), green onions, leaf lettuce, broccoli, cabbage, cauliflower (and other head and stem brassica vegetables), mustard greens (and other leafy brassica greens), spinach, celery, rhubarb (and other leaf petioles), cranberry, cucurbits, fruiting vegetables (except tomatoes), non-bearing food crops, and strawberry. Do not apply more than 5 fl. oz/A of Arrow 2EC Herbicide per application to canola, flax, or mustard seed. **Exceeding these recommendations may result in unacceptable crop injury.**

Do not apply under conditions of stress. Applying Arrow 2EC Herbicide under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity, and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate Arrow 2EC Herbicide effectively and will be less susceptible to herbicide activity.

Best perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices (discing, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, such as continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, results in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, no fewer than two Arrow 2EC Herbicide applications per season per year are recommended at the appropriate weed-growth stage rate under continuous no-till conditions.

Grass crops such as corn, rice, sorghum, small grains, or turf, etc. are highly sensitive to Arrow 2EC Herbicide.

While all the vegetable crops on this label have been tested and are tolerant to Arrow 2EC Herbicide, not all specialty varieties of these crops have been tested. It is advised that, before applying Arrow 2EC Herbicide to specialty varieties of vegetable crops on this label, crop tolerance be investigated first using a small section of the field. It is possible that injury symptoms can occur. Symptoms may appear as leaf speckling or stunting.

Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.

Tank mixes of Arrow 2EC Herbicide and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of Arrow 2EC Herbicide may be necessary.

### **AVOID SPRAY DRIFT**

Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:

- Do not spray if wind speeds are or become excessive. Do not spray if wind speed is 10 MPH or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions. Do not spray if winds are gusty.
- Use extreme caution when conditions are favorable for drift (high temperatures, drought, low relative humidity), especially when sensitive plants are located nearby.
- Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.
- Further reductions in drift can be obtained by:
  1. Using large droplet size sprays. Do not use nozzles that produce small droplets. Orient nozzles downward and slightly backward as needed to reduce drift for ground applications.
  2. Orienting nozzles straight back with the wind stream, using straight stream orifices for aerial applications. Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 3 GPA volume. Application height and boom length should be set according to manufacturer's instructions to minimize drift.

- Increasing the volume of spray mixture (for example a minimum of 10 GPA for ground applications) by using higher flow rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift.
- Applying as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

**Do not apply under conditions involving possible drift to food, forage, or other planting that might be damaged or the crops thereof rendered unfit for sale, use, or consumption.**

### PREHARVEST INTERVAL

The required days to wait between the last application and harvest are given in ( ) after each crop name.

### CROP-SPECIFIC RESTRICTIONS AND LIMITATIONS FOR ARROW 2EC HERBICIDE

CROPS (1)	USE RATES PER ACRE	CROP OIL CONCENTRATE RATES PER ACRE (2)	SPECIAL USE INSTRUCTIONS
<b>Alfalfa (including: Sainfoin, Holy Clover, Birdsfoot Trefoil) (3)</b> (15 days before grazing, feeding or harvesting (cutting) for forage or hay)	6-16 fl. oz. (4)	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air (5)	Do not plant rotational crops until 30 days after application of Arrow 2EC Herbicide. (6) The addition of AMS has shown improved grass control for difficult to control species including quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.

<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Beans, Dry</b> (30 days)	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Refer to appropriate Table for reduced rate recommendations for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Beet, Garden</b> (30 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Carrot</b> (30 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Canola</b> (75 days)	4-5 fl. oz.	1% v/v in the finished spray volume	Do not apply after crop has begun bolting. Crop injury may occur when Arrow 2EC Herbicide is applied during the bloom period. Do not exceed 5 fl. oz. of Arrow 2EC Herbicide (0.08 lb. ai) per acre in a season.

*(continued)*



<b>CROPS <sup>(1)</sup></b>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE <sup>(2)</sup></b>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Celery (includes Cardoon, Chinese Celery, Celtuce, Florence Fennel, Swiss Chard)</b> (30 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Clover (Idaho, Oregon, and Washington only)</b> (15 days before grazing, feeding, or harvesting (cutting) for forage or hay)	6-16 fl. oz.	1% v/v in the finished spray volume	For use on clover grown in the states of Idaho, Oregon, and Washington only. In Oregon, for use on clover grown for seed, use 6-8 fl. oz. per acre for annual grass control and 8-10 fl. oz. per acre for perennial grass control. Do not exceed 16 fl. oz. of Arrow 2EC Herbicide (0.25 lb. ai) per acre in a season.
<b>Cotton</b> (60 days)	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Do not graze treated fields or feed treated forage or hay to livestock. The addition of AMS has shown improved grass control for difficult

<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Cotton</b> (60 days) <i>(continued)</i>	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	to control species including quack-grass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Cranberry</b> (30 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Cucurbits (including Cantaloupes [all], Cucumber, Gherkin, Honeydew Melon, Muskmelons [all], Pumpkin, Squash [all], Watermelon)</b> (14 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.

*(continued)*

<b>CROPS <sup>(1)</sup></b>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE <sup>(2)</sup></b>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Fallow Land, Conifer Trees (and other non-producing agricultural areas), Non-Crop or Non-Planted Areas</b>	6-16 fl. oz.	1% v/v (but not less than 1 pt./A) in the finished spray volume using a crop oil concentrate containing at least 15% emulsifier.	Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.
<b>Flax</b> (75 days)	4-5 fl. oz.	1% v/v in the finished spray volume	Apply prior to bloom. Crop injury may occur when Arrow 2EC Herbicide is applied during the bloom period. Do not exceed 5 fl. oz. of Arrow 2EC Herbicide (0.08 lb.ai) per acre in a season.
<b>Fruiting Vegetables</b> ([except Tomato] including Eggplant, Ground-cherry,	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.

*(continued)*

<b>CROPS <sup>(1)</sup></b>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE <sup>(2)</sup></b>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Pepino, Peppers [all], Tomatillo</b> (20 days) <i>(continued)</i>	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Head and Stem Brassica Vegetables (including Broccoli, Cabbage, Cauliflower, Brussels Sprouts)</b> (30 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Leafy Brassica Greens (including Broccoli Raab; Cabbage, Chinese (Bok Choy; Collards; Kale;</b>	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.

<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Mizuna; Mustard Greens, Mustard Spinach, Rape Greens, Turnip Greens)</b> (14 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Leaf Lettuce</b> (14 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Mint</b> (21 days)	6-16 fl. oz. <sup>(4)</sup>	1 qt. by ground or 1.0% v/v (but not less than 1 pt. per acre by air)	Do not apply more than 16 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Mustard Seed</b> (75 days)	4-5 fl. oz.	1% v/v in the finished spray volume	Do not apply after crop has begun bolting. Crop injury may occur when Arrow 2EC Herbicide is applied during the bloom period. Do not exceed 5 fl. oz. of Arrow 2EC Herbicide (0.08 lb. ai) per acre in a season.

*(continued)*

<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Onions (Dry Bulbs Only), Garlic, Shallots (Dry Bulbs Only)</b> (45 days)	6-16 fl. oz. (7), (8)	1% v/v in the finished spray volume	Minimum of 20 gals./A spray volume by ground in entire U.S.
<b>Onions, Green including Leeks, Scallions, or Spring Onions, Japanese Bunching Onions, Green Shallots, Green Eschalots</b> (14 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Ornamentals</b>	6-16 fl. oz.	Use of crop oil concentrate is not recommended since it may	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25%v/v). Sugar maples cannot be tapped
<b>Non-Bearing Food Crops</b>	6-8 fl. oz. <sup>(8)</sup>		

*(continued)*

<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Non-Bearing Food Crops</b> <i>(continued)</i>	6-8 fl. oz. <sup>(8)</sup>	injure flowers and foliage. See Special use Instructions	for syrup within one year of Arrow 2EC Herbicide application.
<b>Peanuts</b> (40 days)	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	The addition of AMS has shown improved grass control for difficult to control species including quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Potato</b> (30 days)	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	The addition of AMS has shown improved grass control for difficult to control species including quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Radish</b> (15 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. Do not apply more than 16 fl. oz. (0.25 lb. ai) per acre in a season. For repeat applications make on a minimum of a 14-day interval.
<b>Root Vegetables (except Radish)</b>	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.

<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>including Chicory, Ginseng, Horseradish, Turnip</b> (30 days) <i>(continued)</i>	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Rhubarb</b> (30 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Soybean</b> (60 days)	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Do not graze treated fields or feed treated forage or hay to livestock. Refer to appropriate Table for reduced rate recommendations for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Spinach</b> (14 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.



<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>Strawberry</b> (4 days)	6-8 fl. oz.	1% v/v in the finished spray volume	Do not apply more than 8 fl. oz./A in a single application. For repeat applications make on a minimum of a 14-day interval.
<b>Sugar Beet</b> (40 days)	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	Refer to appropriate table for reduced rate recommendations for the control of small annual grasses. The addition of AMS has shown improved grass control for difficult to control species including quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Sunflower</b> (70 days)	6-16 fl. oz.	1 qt. by ground or 1.0% v/v (but not less than 1 pt./A) by air <sup>(5)</sup>	The addition of AMS has shown improved grass control for difficult to control species including quackgrass, Rhizome johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
<b>Sweet Potato, Yam, and other tuberous and corm vegetables</b>	6-16 fl. oz.	1% v/v in the finished spray volume	

<b>CROPS</b> <sup>(1)</sup>	<b>USE RATES PER ACRE</b>	<b>CROP OIL CONCENTRATE RATES PER ACRE</b> <sup>(2)</sup>	<b>SPECIAL USE INSTRUCTIONS</b>
<b>(except potato) including Artichoke (Chinese, Jerusalem), Cassava (Bitter, Sweet), Ginger (30 days)</b> <i>(continued)</i>	6-16 fl. oz.	1% v/v in the finished spray volume	
<b>Tomato (20 days)</b>	6-16 fl. oz.	1% v/v in the finished spray volume	

N/A= Not Applicable

<sup>(1)</sup> Arrow 2EC Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>(2)</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality, and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the ADDITION OF ADJUVANT AND CROP OIL CONCENTRATE section for further information.

<sup>(3)</sup> Arrow 2EC Herbicide may be applied to seedling or established alfalfa grown for seed, hay, silage, green chop, or direct grazing.

- (4) For weed control in established alfalfa and mint, the minimum use rate is 8 fl. oz./A.
- (5) 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to Arrow 2EC Herbicide applications in addition to the recommended rate of crop oil concentrate.
- (6) Do not apply Arrow 2EC Herbicide and 2,4-DB as a tank mix to alfalfa unless the 60 day feeding, grazing, and harvesting restriction on the 2,4-DB label can be observed.
- (7) For ground applications to garlic or shallots, do not exceed 8 fl. oz./A in a single application. For air applications to onions, garlic, or shallots, do not exceed 8 fl. oz./A in a single application. For garlic and shallots, do not exceed 2 applications per season.
- (8) If Arrow 2EC Herbicide is applied as a spot treatment to onions, garlic, shallots, or non-bearing food crops, care should be taken to not exceed the maximum rate allowed on a PER ACRE basis or crop injury may occur.

**DIRECTIONS FOR USE IN SOYBEANS, COTTON, SUGAR BEETS, ONIONS (Dry Bulbs and Green), GARLIC, SHALLOTS (Dry Bulbs and Green), ALFALFA, PEANUTS, DRY BEANS, SUNFLOWER, CANOLA, FLAX, MUSTARD SEED, POTATO, SWEET POTATO, YAM (and other Tuberous\* and Corm\* Vegetables), TOMATOES, PEPPERS (bell and non-bell), EGGPLANTS (and other Fruiting Vegetables), CARROT, RADISH, GARDEN BEET, HORSERADISH (and other Root Vegetables\*\*), LEAF LETTUCE, BROCCOLI, CABBAGE, CAULIFLOWER (and other Head and Stem Brassica Vegetables\*\*\*), MUSTARD GREENS (and other leafy brassica greens\*\*\*\*), SPINACH, CELERY, RHUBARB (and other Leaf Petioles\*\*\*\*), CRANBERRY, STRAWBERRY, SQUASH (including PUMPKINS), CUCUMBER, MELONS (including CANTALOUPEs and WATERMELONS), MINT, and CLOVER (grown in Idaho, Oregon, and Washington only), CONIFER TREES, NON-BEARING FOOD CROPS, AND NON-CROP or NON-PLANTED AREAS.**

\*Other tuber and corm vegetables approved for use with Arrow 2EC Herbicide include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible burdock, edible canna, bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, tanier, turmeric, and bean yam

\*\*Other root vegetables approved for use with Arrow 2EC Herbicide include: burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret and turnip

\*\*\*Other head and stem brassica vegetables approved include: Chinese broccoli; Brussels sprouts; Chinese (napa) cabbage; Chinese mustard; cavalo broccolo; and kohlrabi

\*\*\*\*Other leaf petiole crops include: cardoon, Chinese celery, celtuce, Florence fennel, and Swiss chard

\*\*\*\*\*Other leafy brassica greens approved for use with Arrow 2EC Herbicide include: broccoli raab; cabbage, Chinese (Bok Choy); collards; kale; mizuna; mustard greens; mustard spinach; rape greens; turnip greens.

**IMPORTANT:** Plant tolerance to Arrow 2EC Herbicide at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of Arrow 2EC Herbicide have investigated the safety factor to plants not listed on the label.

## **NON-BEARING FOOD CROPS**

**Arrow 2EC Herbicide SHOULD NOT BE APPLIED TO NON-BEARING FRUIT OR NUT CROPS WHICH ARE GROWN FOR ROOT STOCK.**

**Crop injury to non-bearing fruit and nut crops can occur if Arrow 2EC Herbicide is improperly applied. Arrow 2EC Herbicide should not be applied directly over the top of these plant types. Instead, spray should be directed at the base of the plant where grassy weeds are growing near the ground.**

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following Arrow 2EC Herbicide application.

<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>
Apples	<i>Malus</i> spp.
Berries	<i>Vaccinium</i> spp.
	<i>Rubus</i> spp.
Cherry, Sweet	<i>Prunus avium</i>
Citrus Fruits	<i>Citrus</i> spp.
Grapes	<i>Vitis</i> spp.
Olives	<i>Olea</i> spp.
Peach	<i>Prunus persica</i>
Pears	<i>Pyrus communis</i>
Prunes	<i>Prunus</i> spp.
Stone Fruits	<i>Prunus</i> spp.
Strawberries	<i>Fragaria</i> spp.
<b>Tree Nuts:</b>	
Almonds	<i>Prunus dulcis</i>
Filbert	<i>Corylus maxima</i>
Pecan	<i>Carya illinoensis</i>
Pistachio	<i>Pistacia vera</i>
Walnut	<i>Juglans</i> spp.

## CONIFER TREES

Arrow 2EC Herbicide can be used to control labeled grasses in Christmas tree farms, conifer nurseries, and conifer plantations (but not in forests).

COMMON NAME	SCIENTIFIC NAME
Arborvitae, American	<i>Thuja occidentalis</i>
Cedars	<i>Cedrus</i> spp.
Cypress	<i>Taxodium</i> spp.
Douglas Fir	<i>Pseudotsuga menziesii</i>
Firs	<i>Abies</i> spp.
Hemlock, Canadian	<i>Tsuga canadensis</i>
Hemlock, Western	<i>Tsuga heterophylla</i>
Pines	<i>Pinus</i> spp.
Spruces	<i>Picea</i> spp.
Yew	<i>Taxus</i> spp.

## NON-CROP OR NON-PLANTED AREAS

The following areas are considered non-crop or non-planted areas: Rights-of-way including railroads, highways, roads, dividers, medians, pipelines, public utility lines, pumping stations, transformer stations and substations, around airports, electric utilities, commercial buildings, manufacturing plants, storage yards, rail yards, fence lines, parkways, and post-harvest croplands, also beneath greenhouse benches and around golf courses.

## RECOMMENDATIONS FOR ANNUAL GRASSES (EXCEPT IN ESTABLISHED ALFALFA AND MINT)

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

- Do not apply more than 8 fl.oz./A of Arrow 2EC Herbicide per application to the following crops: garden beets, carrots, radish (and other root vegetables), green onions, leaf lettuce, broccoli, cabbage, cauliflower (and other head and stem brassica vegetables), mustard greens (and other leafy brassica greens), spinach, celery, rhubarb (and other leaf petioles), cranberry, cucurbits, fruiting vegetables (except tomatoes), non-bearing food crops, and strawberry. Do not apply more than 5 fl. oz./A of Arrow 2EC Herbicide per application to canola, flax, or mustard seed.

<b>GRASS SPECIES</b>	<b>SCIENTIFIC NAME</b>	<b>WEED HEIGHT* INCHES</b>	<b>RATE FL. OZ./ACRE</b>	<b>HIGH RATE (4)</b>
BARNYARDGRASS	<i>Echinochloa crus-galli</i>	2 to 8	6	8
BROADLEAF SIGNALGRASS	<i>Brachiaria platyphylla</i>	2 to 6	6	8
BROME				
California	<i>Bromus carinatus</i>	2 to 6	6	8
Cheatgrass	<i>Bromus secalinus</i>	2 to 6	6	8
Downy	<i>Bromus tectorum</i>	2 to 6	6	8
Ripgut	<i>Bromus diandrus</i>	2 to 6	6	8
CANARYGRASS	<i>Phalaris canariensis</i>	1 to 4	6	8
CRABGRASS				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	6	8
Large	<i>Digitaria sanguinalis</i>	2 to 6**	6	8
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	6	8
Southern	<i>Digitaria ciliaris</i>	2 to 6**	6	8
CROWFOOTGRASS	<i>Dactyloctenium aegyptium</i>	2 to 6**	6	8
FALL PANICUM	<i>Panicum dichotomiflorum</i>	2 to 8	6	8

<b>GRASS SPECIES</b>	<b>SCIENTIFIC NAME</b>	<b>WEED HEIGHT* INCHES</b>	<b>RATE FL. OZ./ACRE</b>	<b>HIGH RATE (4)</b>
FIELD SANDBUR	<i>Cenchrus incertus</i>	2 to 6	6	8
FOXTAIL				
Giant	<i>Setaria faberi</i>	2 to 12	6	8
Green	<i>Setaria viridis</i>	2 to 8	6	8
Yellow	<i>Setaria glauca</i>	2 to 8	6	8
GOOSEGRASS	<i>Eleusine indica</i>	2 to 6**	6	8
ITCHGRASS	<i>Rottboellia exaltata</i>	2 to 6	6	8
JUNGLERICE	<i>Echinochloa colona</i>	2 to 6	6	8
LOVEGRASS (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	6	8
RABBITSFOOTGRASS	<i>Polypogon monspeliensis</i>	1 to 4	6	8
RED RICE	<i>Oryza sativa</i>	1 to 3	6	8
RYEGRASS				
Hardy	<i>Lolium remotum</i>	2 to 6	6	8
Italian	<i>Lolium multiflorum</i>	2 to 6	6	8
SEEDLING JOHNSONGRASS	<i>Sorghum halepense</i>	4 to 10	6	8
SHATTERCANE	<i>Sorghum bicolor</i>	6 to 18	6	8
SOUTHWESTERN CUPGRASS	<i>Eriochloa gracilis</i>	2 to 6	6	8
SPRANGLETOP				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	6	8
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	6	8



<b>GRASS SPECIES</b>	<b>SCIENTIFIC NAME</b>	<b>WEED HEIGHT* INCHES</b>	<b>RATE FL. OZ./ACRE</b>	<b>HIGH RATE<sup>(4)</sup></b>
Mexican	<i>Leptochloa uninervia</i>	2 to 6	6	8
Red	<i>Leptochloa filiformis</i>	2 to 6	6	8
TEXAS PANICUM	<i>Panicum texanum</i>	2 to 6	6	8
VOLUNTEER CEREALS <sup>(3)</sup>				
Barley	<i>Hordeum vulgare</i>	2 to 6	6	8
Oats	<i>Avena sativa</i>	2 to 6	6	8
Rye	<i>Secale cereale</i>	2 to 6	6	8
Wheat	<i>Triticum aestivum</i>	2 to 6	6	8
VOLUNTEER CORN <sup>(2)</sup>	<i>Zea mays</i>	4 to 12	4	6
VOLUNTEER CORN (S. R.) <sup>(1)</sup>	<i>Zea mays</i>	4 to 12	8 (suppression only)	
VOLUNTEER CORN <sup>(2)</sup>	<i>Zea mays</i>	12 to 24	6	8
VOLUNTEER GRAIN SORGHUM	<i>Sorghum bicolor</i>	8 to 12	6	8
WILD OATS	<i>Avena fatua</i>	2 to 6	6	8
WILD PROSO MILLET	<i>Panicum miliaceum</i>	2 to 10	6	8
WITCHGRASS	<i>Panicum capillare</i>	2 to 8	6	8
WOOLLY CUPGRASS	<i>Eriochloa villosa</i>	2 to 8	6	8

\*Generally occurs between 3-leaf stage and tillering

\*\*Length of lateral growth

(1) Sethoxydim-resistant volunteer corn.

(2) Includes Roundup Ready®, Liberty Link®, and IMI-CORN® VOLUNTEER CORN.

(3) When a cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum Arrow 2EC Herbicide use rate for control is 8 fl. oz./A.

(4) Rates higher than 8 fl. oz./A may be applied in certain geographic areas, cropping situations, or environmental conditions where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 8 to 16 fl. oz./A may be applied. Do not apply more than 8 fl. oz./A of Arrow 2EC Herbicide per application to the following crops: garden beets, carrots, radish (and other root vegetables), green onions, leaf lettuce, broccoli, cabbage, cauliflower (and other head and stem brassica vegetables), mustard greens (and other leafy brassica greens), spinach, celery, rhubarb (and other leaf petioles), cranberry, cucurbits, fruiting vegetables (except tomatoes), non-bearing food crops, and strawberry. Do not apply more than 5 fl. oz./A of Arrow 2EC Herbicide per application to canola, flax, or mustard seed.

### **RECOMMENDATIONS FOR ANNUAL & PERENNIAL GRASS CONTROL IN ESTABLISHED ALFALFA AND MINT WITH ARROW 2EC HERBICIDE**

<b>GRASS SPECIES</b>	<b>WEED STAGE</b>	<b>RATE FL. OZ. ACRE</b>	<b>HIGH RATE</b>
Annual & Perennial Grasses Listed in Grass Table	See Table	8	16

**Mowing:** The best control of annual grasses can be achieved by applying Arrow 2EC Herbicide before grass weeds are mowed. Once a grass is mowed it becomes tougher to control as much of the available leaf surface has been removed. In areas without a killing frost, some annuals can over-winter after having been mowed multiple times. These grasses form large crowns and may contain many viable buds. These grasses, even though they may be an annual grass, may require repeated applications of Arrow 2EC Herbicide for partial or complete control.

**Irrigated Alfalfa and Mint:** Irrigation practices can be very critical to the successful use of Arrow 2EC Herbicide in established alfalfa and mint and may be necessary to initiate active growth of the weeds prior to application. Gener-

*(continued)*

ally applications 2 to 4 days after an irrigation are most effective. Irrigation made shortly after application (2 days) can be effective, but more consistent grass control occurs when the irrigation is made before the application.

**Annual Grass Control:** Apply Arrow 2EC Herbicide at the grass sizes indicated in the Recommendation for Annual Grass Table and rates indicated above (8 to 16 fl. oz./A). If a grass has been cut, apply Arrow 2EC Herbicide after active growth has resumed and regrowth has reached the minimum height and before it reaches the maximum height indicated. Apply before the alfalfa/mint canopy covers the grasses and interferes with the spray coverage. Some annual grasses are spring- and summer-germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to clethodim may vary from region to region.

Also some annuals germinate over an extended period of time, and because control of small grasses is desired, applications after each weed flush may be required. As a general rule, spray spring- and summer-germinating grasses as early in the season as possible, after initial green-up. Spray fall-germinating weeds in the fall soon after they begin growing but before any damage is done due to frost. Late fall applications may be less effective due to environmental conditions such as frost, slower plant growth, or the onset of flowering.

**Perennial Grass Control:** Arrow 2EC Herbicide effectively controls perennial grasses such as bermudagrass, johnsongrass, quackgrass, wirestem muhly, tall fescue, foxtail barley, and orchardgrass. Due in part to lack of tillage, perennial grasses are more difficult to control in a perennial crop such as established alfalfa or mint. A program of repeated applications is usually necessary for best results. The best way to control perennial grasses is to do so in the year of stand establishment before rhizomes and stolons become large and difficult to kill.

Use the high rate under heavy grass pressure and/or when grasses are at or near maximum height.

Always add a crop oil concentrate at 1 qt./A by ground or 1% v/v (but not less than 1 pt./A) to the finished spray volume by air.

## RECOMMENDATIONS FOR ANNUAL BLUEGRASS CONTROL WITH ARROW 2EC HERBICIDE

GRASS SPECIES	WEED STAGE	RATE FL. OZ. ACRE	HIGH RATE
ANNUAL BLUEGRASS <i>(Poa annua)</i>	to 4-leaf	6*	16
<p>Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).</p> <p>Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.</p> <p>Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.</p> <p>Always add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.</p>			

\* Use a minimum of 8 fl. oz./A to control annual bluegrass in seedling and established alfalfa and mint.

### DIRECTIONS FOR USE IN DRY BEANS, SOYBEANS, AND SUGAR BEETS AT A REDUCED RATE RECOMMENDATIONS FOR SMALL ANNUAL GRASSES (REDUCED RATE RECOMMENDATIONS NOT FOR USE IN CALIFORNIA)

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low temperatures, and/or under very low humidity.

<b>GRASS SPECIES</b>	<b>SCIENTIFIC NAME</b>	<b>WEED HEIGHT INCHES</b>	<b>RATE FL. OZ./ACRE (1)</b>
BARNYARDGRASS	<i>Echinochloa crus-galli</i>	1 to 4	4
BROADLEAF SIGNALGRASS	<i>Brachiaria platyphylla</i>	1 to 4	5
CRABGRASS			
Large	<i>Digitaria sanguinalis</i>	1 to 3*	4
Large	<i>Digitaria sanguinalis</i>	1 to 4*	5
Smooth	<i>Digitaria ischaemum</i>	1 to 3*	4
Smooth	<i>Digitaria ischaemum</i>	1 to 4*	5
Southern	<i>Digitaria ciliaris</i>	1 to 4*	5
FALL PANICUM	<i>Panicum dichotomiflorum</i>	1 to 4	4
FOXTAIL			
Giant	<i>Setaria faberi</i>	1 to 4	4
Green	<i>Setaria viridis</i>	1 to 4	4
Millet	<i>Setaria italica</i>	1 to 4	5
Yellow	<i>Setaria glauca</i>	1 to 4	4
SEEDLING JOHNSONGRASS	<i>Sorghum halepense</i>	1 to 6	5
SHATTERCANE	<i>Sorghum bicolor</i>	4 to 10	4
TEXAS PANICUM	<i>Panicum texanum</i>	1 to 4	5
VOLUNTEER CORN**	<i>Zea mays</i>	4 to 12	4
WILD PROSO MILLET	<i>Panicum miliaceum</i>	1 to 6	4
WILD OATS	<i>Avena fatua</i>	1 to 4	5

\*Length of lateral growth

\*\*Not S. R. Corn

(1) Always add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.

## DIRECTIONS FOR USE IN CANOLA, FLAX, AND MUSTARD SEED AT REDUCED RATE RECOMMENDATIONS FOR ANNUAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low temperatures, and/or under very low humidity.

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT INCHES	RATE FL. OZ./ACRE
BARNYARDGRASS	<i>Echinochloa crus-galli</i>	1 to 4	4
CRABGRASS			
Large	<i>Digitaria sanguinalis</i>	1 to 4*	5
Smooth	<i>Digitaria ischaemum</i>	1 to 4*	5
FALL PANICUM	<i>Panicum dichotomiflorum</i>	1 to 4	4
FOXTAIL			
Giant	<i>Setaria faberi</i>	1 to 4	4
Green	<i>Setaria viridis</i>	1 to 4	5
Yellow	<i>Setaria glauca</i>	1 to 4	5
SHATTERCANE	<i>Sorghum bicolor</i>	4 to 10	4
VOLUNTEER CEREALS			
Barley	<i>Hordeum vulgare</i>	1 to 4	5
Oats	<i>Avena sativa</i>	1 to 4	5
Wheat	<i>Triticum aestivum</i>	1 to 4	5
VOLUNTEER CORN**	<i>Zea mays</i>	4 to 12	4
WILD OATS	<i>Avena fatua</i>	1 to 4	5
WILD PROSO MILLET	<i>Panicum miliaceum</i>	1 to 6	4

\*Length of lateral growth

\*\*Not S. R. Corn

## RECOMMENDATIONS FOR PERENNIAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- Do not apply more than 8 fl. oz./A of Arrow 2EC Herbicide per application to the following crops: garden beets, carrots, radish (and other root vegetables), green onions, leaf lettuce, broccoli, cabbage, cauliflower (and other head and stem brassica vegetables), mustard greens (and other leafy brassica greens), spinach, celery, rhubarb (and other leaf petioles), cranberry, cucurbits, fruiting vegetables (except tomatoes), non-bearing food crops, and strawberry. Do not apply more than 5 fl. oz./A of Arrow 2EC Herbicide per application to canola, flax, or mustard seed.

GRASS SPECIES	WEED HEIGHT INCHES	RATE FL. OZ./ACRE	HIGH RATE
<b>Bermudagrass (<i>Cynodon dactylon</i>)</b>			
First Application	3 (or up to 6" runners)	8	16
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	8	16
<b>Fescue, tall (<i>Festuca arundinacea</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16
<b>Foxtail Barley (<i>Hordeum jubatum</i>)</b>			
First Application	2 to 6	8	16
Repeat Application(s) (if regrowth occurs)	2 to 6	8	16

*(continued)*

GRASS SPECIES	WEED HEIGHT INCHES	RATE FL. OZ./ACRE	HIGH RATE
<b>Orchardgrass (<i>Dactylis glomerata</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16
<b>Quackgrass* (<i>Agropyron repens</i>)</b>			
First Application	4 to 12	8	16
Repeat Application(s) (if regrowth occurs)	4 to 12	8	16
<b>Rhizome Johnsongrass (<i>Sorghum halepense</i>)</b>			
First Application	12 to 24	8	16
Repeat Application(s) (if regrowth occurs)	6 to 18	6	8
<b>Wirestem Muhly (<i>Muhlenbergia frondosa</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16
<b>Perennial Bluegrass* [Roughstalk (<i>Poa trivialis</i>)] [Kentucky (<i>Poa pratensis</i>)]</b>			
First Application	2 to 4	8	16
Repeat Application(s)	2 to 4	8	16

\*Control of quackgrass and perennial bluegrass with Arrow 2EC Herbicide may be enhanced by adding AMS at 2.5 to 4.0 lbs./A.



## **TANK MIXES GENERAL INFORMATION**

The labels for each of the herbicides recommended for tank mixing with Arrow 2EC Herbicide are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the Arrow 2EC Herbicide label in certain considerations. Those concerns may include, but are not limited to:

1. Geographic restrictions-all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (e.g. protective clothing, reentry time, posting);
5. Soil type or soil characteristics (e.g. pH, OM);
6. Maximum dosage or number of applications per season;
7. Rain free period required; or
8. Application timing (e.g. pre-harvest interval)

9. **FOR ALL CROPS EXCEPT CANOLA, CLOVER, FLAX, MUSTARD, AND RADISH:** DO NOT EXCEED A TOTAL OF 32 FL. OZ./A (0.5 LB./AI/A) OF ARROW 2EC HERBICIDE PER SEASON, WHETHER APPLIED ALONE OR IN TANK MIX APPLICATIONS.

**FOR CLOVER AND RADISH:** DO NOT EXCEED A TOTAL OF 16 FL. OZ. ARROW 2EC HERBICIDE (0.25 LB. OF AI) PER ACRE PER SEASON.

**FOR CANOLA, MUSTARD, AND FLAX:** DO NOT EXCEED A TOTAL OF 5 FL. OZ. ARROW 2EC HERBICIDE (0.08 LB. OF AI) PER ACRE PER SEASON.

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

### **TANK MIX APPLICATION OF ARROW 2EC HERBICIDE AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS**

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label.

- Apply when the first grass or broadleaf weed species in a mixed population reaches the recommended height or growth stage for treatment.
- Apply under favorable soil moisture and humidity, which exist a few days after rainfall or within seven days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate recommended for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide, as specified in the respective size and rate tables.
- Do not tank mix Arrow 2EC Herbicide when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

### **MIXING INSTRUCTIONS**

Maintain agitation throughout the spray application. Failure to agitate the spray volume may result in improper mixing of the herbicides and unsatisfactory weed control. Mixing and compatibility qualities should be verified by a jar test.

Arrow 2EC Herbicide Tank Mix: Add 1/2 of the required water to the spray tank and begin agitation. Add the required amount of Arrow 2EC Herbicide and mix thoroughly. Then add the required amount of tank mix partner and continue mixing. Finally add the required amount of crop oil concentrate and/or the nitrogen fertilizer and the remaining water.

### **INFORMATION ON ANTAGONISM**

Tank mixes of Arrow 2EC Herbicide with postemergence broadleaf herbicides have shown some reduction or failure to control certain grass species, which would have otherwise been controlled when Arrow 2EC Herbicide is applied alone. Activity of the postemergence broadleaf herbicide in the tank mix is not affected.

**Table 1****ARROW 2EC HERBICIDE TANK MIXES<sup>(3)</sup> TO CONTROL ANNUAL GRASSES WHEN USED AS A BURNDOWN IN NO-TILL SOYBEANS**

<b>PRODUCT</b>	<b>PRODUCT ACRE RATE (1)</b>	<b>GRASS HEIGHT</b>	<b>CROP OIL CONC. (2)</b>	<b>28%N OR 32%N QTS./A OR 2.5 TO 4.0 LBS. AMS</b>
Arrow 2EC  +  2,4-D Ester*	3 fl. oz.	Foxtail 1 to 3" Fall Panicum 1 to 3"	1 qt./A	1 to 2 qts./A <b>or</b> 2.5 to 4.0 lbs. AMS
	4 fl. oz.	Foxtail 1 to 4" Fall Panicum 1 to 4"	1 qt./A	1 to 2 qts./A <b>or</b> 2.5 to 4.0 lbs. AMS
	6 to 8 fl. oz. + 0.5 lb. ai	(See Grass Chart for grasses claimed)	1 qt./A	1 to 2 qts./A <b>or</b> 2.5 to 4.0 lbs. AMS

\*2,4-D Ester should not be used where drift-sensitive crops may be grown.

- (1) If regrowth occurs or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide according to the appropriate size and rate recommendations.
- (2) Always use a crop oil concentrate at the listed rate in the finished spray volume.
- (3) The following products can be tank mixed with Arrow 2EC Herbicide plus 2,4-D Ester: Dual 8E®, Dual II®, Dual Magnum®, Prowl®, Sencor®, and Sencor plus the Dual products.

Table 2

**ARROW 2EC HERBICIDE TANK MIXES  
WITH BROADLEAF HERBICIDES FOR SOYBEANS  
(REFER TO THE RECOMMENDATION TABLES ABOVE  
FOR SPECIFIC GRASSES AND GROWTH STAGES)**

PRODUCT (2)	APPLICATION RATES/ACRE (1)			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONC. (3) (V/V)	
			GROUND	AIR
Arrow 2EC + Cobra® Herbicide	6 to 8 fl. oz. + 12.5 fl. oz.	8 to 16 fl. oz. + 12.5 fl. oz.	0.5 to 1%	1%
Arrow 2EC + Basagran® 4 SL	8 to 10 fl. oz. + 1 to 2 pts.	10 to 16 fl. oz. + 1 to 2 pts.	1%	1%
Arrow 2EC + Blazer® 2 SL	6 to 8 fl. oz. + 1 to 1.5 pts.	6 to 8 fl. oz. + 1 to 1.5 pts.	0.5 to 1%	1%
Arrow 2EC + Flexstar® HL Herbicide (5)	6 to 8 fl. oz. Refer to the Flexstar HL Herbicide label for specific application rates	6 to 8 fl. oz. Refer to the Flexstar HL Herbicide label for specific application rates	1%	1%
Arrow 2EC + Classic® 25 DG	8 to 10 fl. oz. + 0.5 to 0.75 oz.	10 to 16 fl. oz. + 0.5 to 0.75 oz.	1%	1%
Arrow 2EC (4) + Pursuit®	6 to 8 fl. oz. + 4 fl. oz.	8 to 16 fl. oz. + 4 fl. oz.	1%	1%

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONC. <sup>(3)</sup> (V/V)	
			GROUND	AIR
Arrow 2EC <sup>(4)</sup> + Reflex® 2 LC	6 to 8 fl. oz. + 0.75 to 1.5 pts.	8 to 16 fl. oz. + 0.75 to 1.5 pts.	0.5% to 1%	1%
Arrow 2EC <sup>(4)</sup> + Galaxy™	8 to 10 fl. oz. + 32 fl. oz.	–	0.5%	1%
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + Classic 25 DG	8 to 10 fl. oz. + 6 to 8 fl. oz. + 0.5 to 0.75 oz.	–	0.5%	1%
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + Basagran 4 SL	8 to 10 fl. oz. + 6 to 10 fl. oz. + 1 to 1.5 pts.	–	0.5%	1%
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + Pursuit	8 to 10 fl. oz. + 6 to 10 fl. oz. + 4 fl. oz.	–	0.5%	1%
Arrow 2EC <sup>(4)</sup> + Storm®	8 to 10 fl. oz. + 1.5 pts.	–	0.5%	1%

*(continued)*

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONC. <sup>(3)</sup> (V/V)	
			GROUND	AIR
Arrow 2EC <sup>(4)</sup> + Resource <sup>®</sup> Herbicide + Pursuit	8 to 10 fl. oz. + 4 fl. oz. + 4 fl. oz.	—	1%	1%
Arrow 2EC <sup>(4)</sup> + Resource Herbicide + Basagran Herbicide	8 to 10 fl. oz. + 4 fl. oz. + 1 pt.	—	1%	1%
Arrow 2EC <sup>(4)</sup> + Resource Herbicide + Classic Herbicide	8 to 10 fl. oz. + 4 fl. oz. + 0.5 fl. oz.	—	1%	1%
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + Resource Herbicide	6 to 8 fl. oz. + 6 fl. oz. + 4 fl. oz.	—	0.5%	1%
Arrow 2EC <sup>(4)</sup> + FirstRate <sup>®</sup>	6 to 8 fl. oz. + 0.3 oz.	8 to 16 fl. oz. + 0.3 oz.	1%	—

*(continued)*

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONC. <sup>(3)</sup> (V/V)	
			GROUND	AIR
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + FirstRate	6 to 8 fl. oz. + 6 to 8 fl. oz. + 0.3 oz.	8 to 16 fl. oz. + 6 to 8 fl. oz. + 0.3 oz.	1%	–
Arrow 2EC <sup>(4)</sup> + Raptor® (1 AS)	6 to 8 fl. oz. + 4 to 5 fl. oz.	–	1%	–
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + Raptor (1 AS)	6 to 8 fl. oz. + 6 to 8 fl. oz. + 4 to 5 fl. oz.	–	1%	–
Arrow 2EC <sup>(4)</sup> + Synchrony® STS™	6 to 8 fl. oz. <sup>(6)</sup> + 0.5 oz.	–	1 qt./A	–
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + Synchrony STS	6 to 8 fl. oz. <sup>(6)</sup> + 4 to 8 fl. oz. + 0.5 oz.	–	1 pt./A	–
Arrow 2EC <sup>(4)</sup> + Resource Herbicide	6 to 8 fl. oz. + 4 to 12 fl. oz.	–	1 qt./A	–

*(continued)*

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>			
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONC. <sup>(3)</sup> (V/V)	
			GROUND	AIR
Arrow 2EC <sup>(4)</sup> + Frontrow™	8 to 10 fl. oz. + Refer to Frontrow label for use rates	–	1%	–

- <sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- <sup>(2)</sup> Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- <sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- <sup>(4)</sup> The addition of 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N) is recommended when Arrow 2EC Herbicide is tank mixed with Pursuit, Resource, Galaxy, Storm, FirstRate, Synchrony, Raptor, Frontrow, Cobra plus Classic, Cobra plus Basagran, Cobra plus Pursuit, Cobra plus FirstRate, Cobra plus Synchrony, and Cobra plus Raptor. An equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.
- <sup>(5)</sup> Refer to Flexstar HL label for geographic and rotational restrictions.
- <sup>(6)</sup> Annual grasses and sizes controlled with these tank mixtures are those which are identified in the DIRECTIONS FOR USE IN SOYBEANS AT A REDUCED RATE table.



**Table 3****ARROW 2EC HERBICIDE TANK MIXED WITH COBRA HERBICIDE AND MSMA APPLIED POST DIRECTED TO COTTON**

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONC. <sup>(3)</sup>	COMMENTS
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	
Arrow 2EC <sup>(4)</sup> + Cobra Herbicide + MSMA (4.0 lbs./gal.) or MSMA (6.6 lbs./gal.)	6 to 8 fl. oz.	8 to 16 fl. oz.	1% v/v	Reduce broadcast rate in proportion to the band area actually treated.
	See Cobra label for rates to control broadleaf weeds and height limitations for cotton. Refer to the Arrow 2EC Herbicide label for weed height and species controlled.			
	See MSMA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the Arrow 2EC Herbicide label for weed height and species controlled.			

- (1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- (2) Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- (4) If at the time of application, grass height is so tall that post-directed applications cannot get good coverage over the top of the grassy weeds, then poor control may result and a second (non-post directed) application of Arrow 2EC Herbicide may be necessary.

**Table 4**

**ARROW 2EC HERBICIDE TANK MIXED  
WITH BUCTRIL® 4 EC HERBICIDE TO CONTROL EMERGED  
WEEDS IN BXN COTTON AS A BROADCAST APPLICATION**

PRODUCT (2)	APPLICATION RATES/ACRE (1)	CROP OIL CONC. PER ACRE (3)	COMMENTS (7)
	ANNUAL GRASSES		
Arrow 2EC + Buctril 4 EC Herbicide (4,5,6)	8 to 16 fl. oz.	1 qt./A	See charts for grasses controlled.
	See Buctril 4 EC Herbicide label for rates to control broadleaf weeds and height limitations for cotton.		

- (1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide at the recommended rate with the appropriate amount of crop oil concentrate in a non-Buctril tank mix.
- (2) Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage.
- (3) Always add a crop oil concentrate at 1 qt./A by ground in the finished spray solution.
- (4) Applications of Buctril 4 EC can be made only to cotton that has been genetically modified for crop tolerance to postemergence over-the-top applications of bromoxynil.
- (5) Do not apply the Arrow 2EC Herbicide plus Buctril tank mix within 75 days of harvest.
- (6) Do not exceed two applications of Buctril before cotton is 12 inches tall and one application after 12 inches tall.
- (7) Use a minimum of 10 gals. of spray solution per acre.

**Table 5****ARROW 2EC HERBICIDE TANK MIXED WITH  
STINGER® HERBICIDE APPLIED TO SUGAR BEETS**

<b>PRODUCT (2)</b>	<b>APPLICATION RATES/ACRE (1)</b>		<b>CROP OIL CONC. (3)</b>	
	<b>ANNUAL GRASSES</b>	<b>PERENNIAL GRASSES</b>	<b>GROUND</b>	<b>AIR</b>
Arrow 2EC + Stinger Herbicide	6 to 8 fl. oz.	8 to 16 fl. oz.	1% v/v	
	See Stinger Herbicide label for rates. Refer to the Arrow 2EC Herbicide label for weed height and species controlled.			

- (1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide alone (without a tank mix herbicide) according to the appropriate size and rate recommendations.
- (2) Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

Table 6

## ARROW 2EC HERBICIDE TANK MIXED WITH BETAMIX® OR BETANEX® APPLIED TO SUGAR BEETS

PRODUCT <sup>(2)</sup>	WEEDS CONTROLLED		WEED HEIGHT	APPLICATION RATE/ACRE <sup>(1)</sup>
	Common Name	Scientific Name		
Arrow 2EC <sup>(3)</sup> + Betamix or Betanex	Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 3"	8 fl. oz.
	Foxtail	<i>Setaria</i> spp.	1 to 3"	
	Foxtail Millet	<i>Setaria italica</i>	1 to 3"	
	Wild Oats	<i>Avena fatua</i>	1 to 3"	
	Wild Proso Millet	<i>Panicum miliaceum</i>	1 to 3"	
			See Betamix label for rates to control broad-leaf weeds. <b>No additives are recommended in the tank mix.</b>	
			See Betanex label for rates to control broad-leaf weeds. <b>No additives are recommended in the tank mix.</b>	

- (1) Do not use crop oil concentrate. No additives are recommended in the tank mix.
- (2) Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- (3) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide at full label rate with appropriate rate of crop oil concentrate.

## **TANK MIX APPLICATION OF ARROW 2EC HERBICIDE AND 2,4-DB HERBICIDE FOR CONTROL OF GRASSES AND BROADLEAF WEEDS IN ALFALFA**

A tank mix of Arrow 2EC Herbicide plus 2,4-DB (up to 1.0 lb./A) can be used to control grass and broadleaf weeds listed on the two product labels. Include a crop oil concentrate containing at least 15% emulsifier at 1% v/v in the finished spray. Follow rate and other recommendations on the individual herbicide labels when applying this tank mix.

**NOTE:** Arrow 2EC Herbicide plus 2,4-DB may increase the severity of crop injury when tank mixed. Alfalfa plants will generally outgrow this temporary crop injury within a few weeks.

## **TANK MIX APPLICATION OF ARROW 2EC HERBICIDE AND PURSUIT HERBICIDE FOR CONTROL OF GRASSES AND BROADLEAF WEEDS IN ALFALFA**

A tank mix of Arrow 2EC Herbicide plus Pursuit Herbicide or Pursuit DG Herbicide can be used to control annual grass and broadleaf weeds listed on the two product labels. Include a crop oil concentrate at 1% v/v in the finished spray. For annual grass control in alfalfa using Arrow 2EC Herbicide plus Pursuit, use 8 to 16 fl. oz/A of Arrow 2EC Herbicide.

Before using this tank mix, read and understand the Pursuit or Pursuit DG Herbicide labels for geographical restrictions and restrictions regarding alfalfa growth stage and type. Failure to do so can result in crop injury to alfalfa.

Do not feed, graze, or harvest alfalfa for 30 days following an application of Pursuit to alfalfa.

**Table 7**

**TANK MIX APPLICATION OF ARROW 2EC HERBICIDE  
AND INSECTICIDES FOR CONTROL OF GRASS WEEDS  
AND INSECTS IN SOYBEANS, COTTON, AND PEANUTS**

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		
	ANNUAL GRASSES	PERENNIAL GRASSES	CROP OIL CONC. <sup>(3)</sup> V/V
Arrow 2EC + Orthene® 75 SP or Orthene 97 (for use in cotton and peanuts only, not soybeans)	6 to 8 fl. oz. + 0.33 to 1.33 lbs. or 0.25 to 1.0 lb.	8 to 16 fl. oz. + 0.33 to 1.33 lbs. or 0.25 to 1.0 lb.	1%
Arrow 2EC + Orthene 90 S <sup>(4)</sup>	6 to 8 fl. oz. + 0.25 to 1 lb.	8 to 16 fl. oz. + 0.25 to 1 lb.	1%
Arrow 2EC + Danitol® 2.4 EC Spray <sup>(5)</sup> (for use in cotton and peanuts only, not soybeans)	6 to 8 fl. oz. + 10 2/3 to 16 fl. oz	8 to 16 fl. oz. + 10 2/3 to 16 fl. oz.	1%

<sup>(1)</sup> If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide alone (without a tank mix insecticide) according to the appropriate size and rate recommendations.

<sup>(2)</sup> Refer to the Arrow 2EC Herbicide and insecticide label for rates, weeds, and insects controlled.

- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- (4) Insecticide tank mix use with Orthene 90S in soybeans is permitted only in a state having an approved Section 24(c) registration for Orthene 90S use in soybeans.
- (5) Danitol tank mix is labeled for use in cotton and peanuts only, not soybeans.

### **TANK MIX APPLICATION OF ARROW 2EC HERBICIDE AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA**

Arrow 2EC Herbicide can be tank mixed with the following insecticides for use in alfalfa: Baythroid®, dimethoate, Lorsban®, Pounce®, or Warrior®. The Arrow 2EC Herbicide rate should be 6 to 8 fl. oz./A for annual grass control in seedling alfalfa, minimum of 8 fl. oz./A for annual grass control in established alfalfa, and 8 to 16 fl. oz./A for perennial grass control. Crop oil concentrate should be added at the rate of 1.0 to 2.0 pts./A. For the Arrow 2EC Herbicide plus Lorsban tank mix, reduce the adjuvant rate down to 1.0 pt./A when the Lorsban rate is 1.0 pt./A or higher.

Certain insecticides may cause temporary phytotoxic symptoms on alfalfa foliage. Refer to the insecticide label for further information. It is suggested that prior to using any of these insecticide/herbicide tank mixtures, that a small area of the field be treated first and observations for crop injury be made prior to treating the whole field. Applications must be made at a timing which falls within the guidelines of the Arrow 2EC Herbicide label relative to weed sizes and application equipment. For these applications, it is necessary to use application equipment designed for herbicide applications.

### **TANK MIX APPLICATION OF ARROW 2EC HERBICIDE AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN MINT**

Arrow 2EC Herbicide can be tank mixed with the following insecticides for use in mint: Orthene®.

The Arrow 2EC Herbicide rate should be 6-8 fl. oz./A for annual grass control in baby mint, minimum of 8 fl. oz./A for annual grass control in established mint, and

8 to 16 fl. oz./A for perennial grass control. Crop oil concentrate should be added at the rate of 1.0 to 2.0 pts./A.

Certain insecticides may cause temporary phytotoxic symptoms on mint foliage. Refer to the insecticide label for further information. It is suggested that prior to using any insecticide/herbicide tank mixtures, that a small area of the field be treated first and observations for crop injury be made prior to treating the whole field.

Applications must be made at a timing which falls within the guidelines of the Arrow 2EC Herbicide label relative to weed sizes and application equipment. For these applications, it is necessary to use application equipment designed for herbicide applications.

**Table 8**

**RECOMMENDATIONS FOR ROUNDUP READY®  
VOLUNTEER CORN CONTROL IN ROUNDUP READY  
SOYBEANS WITH ARROW 2EC HERBICIDE TANK MIX**

<b>PRODUCT</b>	<b>GRASS SPECIES</b>	<b>WEED STAGES</b>	<b>RATE FL. OZ./ACRE</b>	<b>ADJUVANT RATE (1)</b>
Arrow 2EC + Roundup Ultra® (or glyphosate formulations labeled for RR soybean) (3)	RR Volunteer Corn	up to 12 inches up to 24 inches	4 (1) to 6 (2) 6 to 8 + up to 2 qts./A (See Roundup Ultra label, glyphosate labels for use rates)	AMS 2.5 lbs./A

(1) At the 4 fl. oz./A rate of Arrow 2EC Herbicide, the adjuvant recommendation is 1 pt./A COC plus AMS at 2.5 lbs./A.

(2) Use the high rate under heavy grass pressure and/or when grasses are at or near maximum height.

(3) For glyphosate formulations that do not contain a built-in adjuvant system, the adjuvant recommendation is 1 pt./A COC plus AMS (2.5 lbs./A) or NIS at 0.25% v/v plus AMS at 2.5 lbs./A.



**THE MOST RESTRICTIVE LABELING OF ANY  
PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label.
- Apply under favorable soil moisture and humidity which exist a few days after rainfall or within seven days after irrigation.
- Tank mix applications may sometimes result in reduced grass control. If regrowth occurs or an additional flush of new grass emerges, make a second application of Arrow 2EC Herbicide as specified in the respective size and rate tables.
- Do not tank mix Arrow 2EC Herbicide when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.
- This tank mix may be applied postemergence to Roundup Ready soybeans up through the full flowering stage. Do not apply later than 60 days before harvest.
- Avoid contact with foliage, green stems, or fruit crops, or any desirable plants and trees, other than soybeans with the Roundup Ready gene as severe injury or destruction will result.
- Do not allow the Arrow 2EC Herbicide plus Roundup to mist, drip, drift, or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur such as combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.

**DIRECTIONS FOR USE IN FALLOW LAND**

Arrow 2EC Herbicide may be used to control annual and perennial grasses in land that has been left fallow the previous year and other non-producing agricultural areas. Apply Arrow 2EC Herbicide at 6 to 8 fl. oz./A for annual grasses and 8 to 16 fl. oz./A for perennial grasses. When both grass and broadleaf weeds are the target pest, Arrow 2EC Herbicide may be tank mixed with 2,4-D Ester or Banvel SGF Herbicide for broad spectrum control. When both annual and perennial grasses occur in the same field, use a minimum of 8 fl. oz./A Arrow 2EC Herbicide.

## GENERAL INFORMATION

- Use a minimum spray volume of 5 gals./A for aerial applications and 15 gals./A for ground applications.
- Apply only to actively growing grasses when the first grass reaches the recommended weed height as specified by the RECOMMENDATIONS FOR ANNUAL AND PERENNIAL GRASSES section of this label.
- Annual grasses which emerge after the Arrow 2EC Herbicide application will not be controlled and a second application may be necessary.
- The control of perennial grasses may require more than one application in non-tilled areas.
- Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.
- Do not apply to grasses that have tillered, formed seedheads, or exceeded recommended growth stage.
- Do not use flood jet nozzles.
- Do not apply to drought-stressed grasses.
- Do not mow area for two weeks prior to or after the Arrow 2EC Herbicide application.

## ARROW 2EC HERBICIDE IN TANK MIXES TO CONTROL ANNUAL AND PERENNIAL GRASSES IN FALLOW LAND

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONC. <sup>(2)</sup>	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
Arrow 2EC + 2,4-D Ester or Banvel SGF	6 to 8 fl. oz.	8 to 16 fl. oz.	1% v/v	
	0.5 lb.			
	See Banvel SGF label for rates.			

<sup>(1)</sup> Refer to Arrow 2EC Herbicide label for weed height and species control. Review Banvel SGF Herbicide and 2,4-D labels for crop restrictions, use rates, and weeds controlled.

<sup>(2)</sup> Always use a crop oil concentrate or methylated seed oil containing at least 15% emulsifier at the listed rate (but not less than 1 pt./A) in the finished spray volume.

## ARROW 2EC HERBICIDE FOR THE CONTROL AND/OR SUPPRESSION OF TALL FESCUE IN NATIVE PRAIRIE WARM-SEASON GRASS RESTORATION PROJECTS

PRODUCT	PRODUCT RATE	GRASS WEEDS CONTROLLED/SUPPRESSED		WEED STAGE
		COMMON NAME	SCIENTIFIC NAME	
Arrow 2EC	10 to 12 fl. oz./A	Tall Fescue	<i>Festuca arundinacea</i>	4 to 6 in. (40 to 60% green-up)

**Adjuvant:** Arrow 2EC Herbicide must be applied with crop oil concentrate at 1 qt./A, plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A. Recommended Mixing Order: Thoroughly mix spray grade ammonium sulfate in water, add Arrow 2EC Herbicide, then add crop oil concentrate.

### SPECIAL APPLICATION INSTRUCTIONS/PRECAUTIONS

- Burn or mow fields a minimum of 3 weeks prior to application to remove excess crop residue. Apply in the spring, at 40 to 60% tall fescue green-up, prior to emergence of warm-season grasses. Do not mow area for 2 weeks after the Arrow 2EC Herbicide application.
- Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 psi at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood nozzles.
- Apply only to fields that have warm-season grasses established for two years. Applications of Arrow 2EC Herbicide to emerged warm-season grasses may cause injury. Do not apply to warm-season grasses grown for seed.
- Do not graze treated fields or feed treated forage and/or hay to livestock.
- Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.

**NOTE:** Arrow 2EC Herbicide applications are most effective if applied when average nighttime temperatures are consistently greater than or equal to 47 degrees Fahrenheit.

## ARROW 2EC HERBICIDE FOR THE SUPPRESSION OF TALL FESCUE SEED-HEADS IN NON-PRODUCING AGRICULTURAL AREAS

PRODUCT	PRODUCT RATE	SUPPRESSION	APPLICATION TIMING
Arrow 2EC	1 1/2 to 2 fl. oz./A	Tall Fescue Seed-Heads ( <i>Festuca arundinacea</i> )	(50 to 90% Tall Fescue green-up)

**Adjuvant:** Arrow 2EC Herbicide must be applied with crop oil concentrate at 1 qt./A plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A. **Recommended Mixing Order:** Thoroughly mix spray grade ammonium sulfate in water, add Arrow 2EC Herbicide, then add crop oil concentrate.

### SPECIAL APPLICATION INSTRUCTIONS/PRECAUTIONS

- Apply at 50 to 90% tall fescue green-up.
- Use the higher Arrow 2EC Herbicide rate if less tall fescue green matter is present.
- Do not mow area for 2 weeks after the Arrow 2EC Herbicide application.
- Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 psi at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood nozzles.
- 2,4-D Ester may be added to this tank mix for broadleaf control (see 2,4-D Ester label for weeds controlled).
- Do not graze treated fields or feed treated forage and/or hay to livestock.
- Do not plant any crop for 30 days after application, unless clethodim is registered for use in that crop.

### DIRECTIONS FOR USE IN ORNAMENTALS

For ornamental plant uses, Arrow 2EC Herbicide can be used to control labeled grass weeds in greenhouses, lathhouses, shadehouses, and around outdoor ornamentals, including nurseries, parks, roadside plantings, and structure landscapes.

**IMPORTANT:** Arrow 2EC Herbicide successfully controls weeds in newly transplanted and established non-grassy ornamentals. Plant tolerance to Arrow 2EC Herbicide at labeled rates has been found to be acceptable for the indicated genera

and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of Arrow 2EC Herbicide has investigated the safety factor to ornamental plants not listed on the label.

The following plants have shown a tolerance for Arrow 2EC Herbicide applications:

### ORNAMENTAL TREES

COMMON NAME	SCIENTIFIC NAME
ALDER, RED	<i>Alnus oregona</i>
ASH	<i>Fraxinus</i> spp.
BASSWOOD	<i>Tilia</i> spp.
BIRCH, EUROPEAN WHITE	<i>Betula pendula</i>
BIRCH, RIVER	<i>Betula nigra</i>
BIRCH, WHITE	<i>Betula papyrifera</i>
CRABAPPLE, FLOWERING	<i>Malus halliana</i>
DOGWOOD, FLOWERING	<i>Cornus florida</i>
GOLDEN CHAIN TREE	<i>Laburnum anagyroides</i>
MAPLES	<i>Acer</i> spp.
MULBERRY, WHITE	<i>Morus alba</i>
OAKS	<i>Quercus</i> spp.
OLIVE, WILD	<i>Elaeagnus angustifolia</i>
REDBUD	<i>Cercis canadensis</i>
SWEET GUM	<i>Liquidambar styraciflua</i>

### GARDEN FLOWERS AND PLANTS

COMMON NAME	SCIENTIFIC NAME
AGERATUM	<i>Ageratum</i> spp.
ALYSSUM*	<i>Lobularia maritima</i>

## GARDEN FLOWERS AND PLANTS *(continued)*

COMMON NAME	SCIENTIFIC NAME
ASPARAGUS FERN	<i>Asparagus setaceus</i>
BLEEDING HEART	<i>Dicentra spectabilis</i>
CAST IRON PLANT	<i>Aspidistra elatior</i>
CHRYSANTHEMUM	<i>Chrysanthemum</i> spp.
CINQUEFOIL	<i>Potentilla</i> spp.
COLEUS	<i>Coleus</i> spp.
CORALBELLS	<i>Heuchera sanguinea</i>
CRANESBILL	<i>Geranium</i> spp.
DAHLIA	<i>Dahlia</i> spp.
DAISY, AFRICAN	<i>Osteospermum fruticosum</i>
DAYLILY	<i>Hemerocallis</i> spp.
DUSTY MILLER	<i>Senecio cineraria</i>
EUONYMUS	<i>Euonymus</i> spp.
GAZANIA	<i>Gazania</i> spp.
GERANIUM, HOUSE	<i>Pelargonium hortorum</i>
HEATHER	<i>Cuphea hyssopifolia</i>
HOSTA	<i>Hosta fortunei</i>
IRIS	<i>Iris</i> spp.
JASMINE TOBACCO	<i>Nicotiana glauca</i>
LOOSESTRIFE	<i>Lythrum salicaria</i>
MARIGOLD	<i>Tagetes</i> spp.
PARTRIDGEBERRY	<i>Mitchella repens</i>
PETUNIA*	<i>Petunia hybrida</i>
PHLOX	<i>Phlox</i> spp.
PINKS	<i>Dianthus</i> spp.

## GARDEN FLOWERS AND PLANTS *(continued)*

COMMON NAME	SCIENTIFIC NAME
PORTULACA	<i>Portulaca grandiflora</i>
SALVIA	<i>Salvia</i> spp.
SAXIFRAGE	<i>Saxifraga</i> spp.
SEDUM	<i>Sedum</i> spp.
SELLOUM	<i>Philodendron selloum</i>
SNAPDRAGON*	<i>Antirrhinum majus</i>
SWEET FLAG	<i>Acorus gramineus</i>
TICKSEED	<i>Coreopsis grandiflora</i>
TOUCH-ME-NOT	<i>Impatiens</i> spp.
VERBENA	<i>Verbena</i> spp.
VIOLET	<i>Viola</i> spp.
YARROW, COMMON	<i>Achillea millefolium</i>
ZINNIA	<i>Zinnia elegans</i>

\*Slight foliage or flower speckling has been observed on these species.

## GROUND COVERS

COMMON NAME	SCIENTIFIC NAME
BUGLEWEED, CARPET	<i>Ajuga reptans</i>
IVY, ENGLISH	<i>Hedera helix</i>
JAPANESE SPURGE	<i>Pachysandra terminalis</i>
LILYTURF	<i>Liriope muscari</i>
MONEYWORT	<i>Lysimachia nummularia</i>
MONDO GRASS, WHITE	<i>Ophiopogon jaburan</i>
MONDO GRASS, DWARF	<i>Ophiopogon japonicus</i>
PERIWINKLE, COMMON	<i>Vinca minor</i>

## SHRUBS

COMMON NAME	SCIENTIFIC NAME
ABELIA	<i>Abelia</i> spp.
ANISE, PURPLE	<i>Illicium floridanum</i>
AUCUBA	<i>Aucuba</i> spp.
AZALEA*	<i>Rhododendron</i> spp.
BAMBOO	<i>Bambusa</i> spp.
BARBERRY, JAPANESE	<i>Berberis thunbergii</i>
BARBERRY, MAGELLAN	<i>Berberis buxifolia</i>
BAYBERRY	<i>Myrica pensylvanica</i>
BOTTLEBRUSH	<i>Callistemon citrinus</i>
BOXWOOD	<i>Buxus sempervirens</i>
CAMELLIA	<i>Camellia japonica</i>
CANDYTUFT	<i>Iberis sempervirens</i>
CLEYERA	<i>Cleyera japonica</i>
CORALBERRY	<i>Ardisia crenata</i>
CRAPE MYRTLE	<i>Lagerstroemia indica</i>
COYOTE BRUSH	<i>Baccharis pilularis</i>
FIG, CREEPING	<i>Ficus pumila</i>
GARDENIA	<i>Gardenia</i> spp.
HOLLY	<i>Ilex</i> spp.
HONEYSUCKLE	<i>Lonicera pileata</i>
INDIAN HAWTHORN	<i>Raphiolepis indica</i>
JASMINE	<i>Jasminum</i> spp.
JASMINE, ASIATIC	<i>Trachelospermum asiaticum</i>
JASMINE, CONFEDERATE	<i>Trachelospermum jasminoides</i>
JUNIPER	<i>Juniperus</i> spp.



## SHRUBS (*continued*)

COMMON NAME	SCIENTIFIC NAME
LANTANA	<i>Lantana</i> spp
NANDINA*	<i>Nandinia domestica</i>
OLEANDER, COMMON	<i>Nerium oleander</i>
OREGON GRAPE	<i>Mahonia aquifolium</i>
PHOTINIA	<i>Photinia</i> spp.
PITTOSPORUM	<i>Pittosporum</i> spp.
PODOCARPUS	<i>Podocarpus</i> spp.
PRIVET	<i>Ligustrum</i> spp.
PYRACANTHA	<i>Pyracantha</i> spp.
RHODODENDRON	<i>Phododendron</i> spp.
ROSE	<i>Rosa</i> spp.
SPIREA	<i>Spiraea bumalda</i>
TEA OLIVE	<i>Osmanthus fragrans</i>
VIBURNUM	<i>Viburnum tinus</i>
WISTERIA	<i>Wisteria</i> spp.
YELLOW SAGE	<i>Lantana camara</i>

\*Slight foliage or flower speckling has been observed on these species.

## RECOMMENDATIONS FOR ANNUAL GRASSES IN ORNAMENTALS

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT INCHES	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
BARNYARDGRASS	<i>Echinochloa crus-galli</i>	2 to 8	8	16
BROADLEAF SIGNALGRASS	<i>Brachiaria platyphylla</i>	2 to 6	8	16
BROME				
California	<i>Bromus carinatus</i>	2 to 6	8	16
Cheatgrass	<i>Bromus secalinus</i>	2 to 6	8	16
Downy	<i>Bromus tectorum</i>	2 to 6	8	16
Ripgut	<i>Bromus diandrus</i>	2 to 6	8	16
CANARYGRASS	<i>Phalaris canariensis</i>	1 to 4	8	16
CRABGRASS				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	8	16
Large	<i>Digitaria sanguinalis</i>	2 to 6**	8	16
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	8	16
Southern	<i>Digitaria ciliaris</i>	2 to 6**	8	16
CROWFOOTGRASS	<i>Dactyloctenium aegyptium</i>	2 to 6**	8	16
FALL PANICUM	<i>Panicum dichotomiflorum</i>	2 to 8	8	16

<b>GRASS SPECIES</b>	<b>SCIENTIFIC NAME</b>	<b>WEED* HEIGHT INCHES</b>	<b>RATE FL. OZ./ACRE (1)</b>	<b>HIGH RATE (2)</b>
FIELD SANDBUR	<i>Cenchrus incertus</i>	2 to 6	8	16
FOXTAIL				
Giant	<i>Setaria faberi</i>	2 to 12	8	16
Green	<i>Setaria viridis</i>	2 to 8	8	16
Yellow	<i>Setaria glauca</i>	2 to 8	8	16
FOXTAIL BARLEY	<i>Hordeum jubatum</i>	2 to 6	8	16
GOOSEGRASS	<i>Eleusine indica</i>	2 to 6**	8	16
ITCHGRASS	<i>Rottboellia exaltata</i>	2 to 6	8	16
JUNGLERICE	<i>Echinochloa colona</i>	2 to 6	8	16
LOVEGRASS (Stinkgrass)	<i>Eragrostis ciliaris</i>	2 to 6	8	16
RABBITSFOOTGRASS	<i>Polypogon monspeliensis</i>	1 to 4	8	16
RED RICE	<i>Oryza sativa</i>	1 to 3	8	16
RYEGRASS				
Hardy	<i>Lolium remotum</i>	2 to 6	8	16
Italian	<i>Lolium multiflorum</i>	2 to 6	8	16
SEEDLING JOHNSONGRASS	<i>Sorghum halepense</i>	4 to 10	8	16
SHATTERCANE	<i>Sorghum bicolor</i>	6 to 18	8	16
SOUTHWESTERN CUPGRASS	<i>Eriochloa gracilis</i>	2 to 6	8	16

(continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT INCHES	RATE FL. OZ./ACRE (1)	HIGH RATE (2)
SPRANGLETOP				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	8	16
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	8	16
Mexican	<i>Leptochloa uninervia</i>	2 to 6	8	16
Red	<i>Leptochloa filiformis</i>	2 to 6	8	16
TEXAS PANICUM	<i>Panicum texanum</i>	2 to 6	8	16
VOLUNTEER CEREALS				
Barley	<i>Hordeum vulgare</i>	2 to 6	8	16
Oats	<i>Avena sativa</i>	2 to 6	8	16
Rye	<i>Secale cereale</i>	2 to 6	8	16
Wheat	<i>Triticum aestivum</i>	2 to 6	8	16
VOLUNTEER CORN	<i>Zea mays</i>	4 to 12	6	8
VOLUNTEER CORN	<i>Zea mays</i>	12 to 24	8	16
VOLUNTEER GRAIN SORGHUM	<i>Sorghum bicolor</i>	8 to 12	8	16
WILD OATS	<i>Avena fatua</i>	2 to 6	8	16
WILD PROSO MILLET	<i>Panicum miliaceum</i>	2 to 10	8	16
WITCHGRASS	<i>Panicum capillare</i>	2 to 8	8	16
WOOLLY CUPGRASS	<i>Eriochloa villosa</i>	2 to 8	8	16

\*Generally occurs between 3-leaf stage and tillering.

\*\*Length of lateral growth.

(1) 8 fl. oz./A=approximately 0.2 fl. oz./1000 sq. ft.

(2) 16 fl. oz./A=approximately 0.4 fl. oz./1000 sq. ft.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

## RECOMMENDATIONS FOR ANNUAL BLUEGRASS CONTROL WITH ARROW 2EC HERBICIDE IN ORNAMENTALS

GRASS SPECIES	WEED STAGE	RATE FL. OZ. ACRE	HIGH RATE
ANNUAL BLUEGRASS ( <i>Poa annua</i> )	to 4-leaf	6	16
<p>Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).</p> <p>Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.</p> <p>Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.</p> <p>Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).</p>			

## RECOMMENDATIONS FOR PERENNIAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

GRASS SPECIES	WEED HEIGHT INCHES	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
<b>Bermudagrass (<i>Cynodon dactylon</i>)</b>			
First Application	3 (or up to 6" runners)	8	16
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	8	16
<b>Quackgrass (<i>Agropyron repens</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16

GRASS SPECIES	WEED HEIGHT INCHES	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
<b>Rhizome Johnsongrass (<i>Sorghum halepense</i>)</b>			
First Application	12 to 24	8	16
Repeat Application(s) (if regrowth occurs)	6 to 18	6	8
<b>Wirestem Muhly (<i>Muhlenbergia frondosa</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16

<sup>(1)</sup> 8 fl. oz./A=approximately 0.2 fl. oz./1000 sq. ft.

<sup>(2)</sup> 16 fl. oz./A=approximately 0.4 fl. oz./1000 sq. ft.

Add non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal or cleaning of equipment. Open dumping is prohibited.

**STORAGE:** Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Do not store diluted spray.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL INFO-TRAC AT (800) 535-5053.**

## **WARRANTY STATEMENT**

MAKHTESHIM-AGAN OF NORTH AMERICA warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of MAKHTESHIM-AGAN OF NORTH AMERICA. All such risks shall be assumed by the Buyer. Except as expressly provided herein, MAKHTESHIM-AGAN OF NORTH AMERICA makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at MAKHTESHIM-AGAN OF NORTH AMERICA's election, the replacement of this product.

BANVEL® – Reg. TM of BASF Corporation  
BASAGRAN® – Reg. TM of BASF AG.  
BAYTHROID® – Reg. TM Bayer Corporation  
BETAMIX® – Reg. TM of The Aventis Group  
BETANEX® – Reg. TM of The Aventis Group  
BLAZER® – Reg. TM of BASF Corporation  
BUCTRIL® – Reg. TM of Rhone-Poulenc  
CLASSIC® – Reg. TM of E.I. duPont de Nermours & Co. Inc.  
COBRA® – Reg. TM of Valent U.S.A. Corporation  
DANITOL® – Reg. TM of Sumitomo Chemical Co., Ltd.  
DUAL®, DUAL II® and DUAL MAGNUM® – Reg. TMs of Novartis Crop Protection, Inc.  
FIRSTRATE® – Reg. TM of Dow AgroSciences LLC  
FLEXSTAR® – Reg. TM of Zeneca Ag. Products  
FRONTROW™ – TM of Dow AgroSciences LLC  
GALAXY® – Reg. TM of BASF AG  
IMI-CORN® – Reg. TM of American Cyanamid Co.  
LIBERTY LINK® – Reg. TM of The Aventis Group  
LORSBAN® – Reg. TM of Dow AgroSciences LLC  
ORTHENE® – Reg. TM of OMS Investments, Inc.  
POUNCE® – Reg. TM FMC Corporation  
PROWL® – Reg. TM of American Cyanamid Co.  
PURSUIT® and PURSUIT DG® – Reg. TMs of American Cyanamid Co.  
RAPTOR® – Reg. TM of American Cyanamid Co.  
REFLEX® – Reg. TM of Zeneca Ag. Products  
RESOURCE® – Reg. TM of Valent U.S.A. Corporation  
ROUNDUP READY® and ROUNDUP ULTRA® – Reg. TMs of Monsanto Company  
SELECT® – Reg. TM of Valent U.S.A. Corporation  
SENCOR® – Reg. TM of Bayer Corporation  
STINGER® – Reg. TM of Dow AgroSciences LLS  
STORM® – Reg. TM of BASF Corporation  
SYNCHRONY® STS™ – Reg. TM and TM of E.I. duPont de Nemours & Col. Inc.  
WARRIOR® – Reg. TM of Zeneca AG. Products





**MAKHTESHIM**  
**A G A N**  
 NORTH AMERICA



# Arrow™ 2EC HERBICIDE

**ACTIVE INGREDIENT % BY WT.**

Clethodim: (E)-2-[1-[(3-chloro-2-propenyl)oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one ..... 26.4%

**OTHER INGREDIENTS:**..... 73.6%

**TOTAL**.....100.0%

Contains petroleum distillate.  
 Contains 2.0 lbs. clethodim per gallon.

**EPA Reg. No. 66222-60**

**EPA Est. No. 5905-GA-001<sup>HC</sup>  
 5905-IA-001<sup>HD</sup>**

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

**FOR EMERGENCY MEDICAL HELP, CALL PROSAR AT 1-877-250-9291, 24 HOURS.**

For additional precautionary, handling and use statements, and First Aid instructions see inside of this booklet.

Letter in lot number corresponds to superscript in EPA Est. No.

This product is not yet registered in California.



**MAKHTESHIM**  
**A G A N**  
 NORTH AMERICA

**Makhteshim Agan of North America, Inc.**  
**MANA**  
 551 Fifth Avenue / Suite 1100  
 New York, NY 10176

**Net Contents: 1 gallon**